

**[0188]** The inlet hole of the microfluidic cartridge, or other microchannel device, is dimensioned in such a way that the droplet of 75 nl can be accurately propelled to the bottom of the inlet hole using, for example, compressed air, or in a manner similar to an inkjet printing method. The microfluidic cartridge is maintained at a temperature above the melting point of the wax thereby permitting the wax to stay in a molten state immediately after it is dispensed. After the drop falls to the bottom of the inlet hole **1901**, the molten wax is drawn into the narrow channel by capillary action, as shown in the sequence of views in FIG. 25B. A shoulder between the inlet hole **1901** and the loading channel can facilitate motion of the TRS. The volume of the narrow section can be designed to be approximately equal to a maximum typical amount that is dispensed into the inlet hole. The narrow section can also be designed so that even though the wax dispensed may vary considerably between a minimum and a maximum shot size, the wax always fills up to, and stops at, the microchannel junction **1907** because the T-junction provides a higher cross section than that of the narrow section and thus reduces the capillary forces.

#### PCR Reagent Mixtures

**[0189]** In various embodiments, the sample for introduction into a lane of the microfluidic cartridge can include a PCR reagent mixture comprising a polymerase enzyme and a plurality of nucleotides.

**[0190]** In various embodiments, preparation of a PCR-ready sample for use with an apparatus and cartridge as described herein can include contacting a neutralized polynucleotide sample with a PCR reagent mixture comprising a polymerase enzyme and a plurality of nucleotides (in some embodiments, the PCR reagent mixture can further include a positive control plasmid and a fluorogenic hybridization probe selective for at least a portion of the plasmid).

**[0191]** The PCR-ready sample can be prepared, for example, using the following steps: (1) collect sample in sample collection buffer, (2) transfer entire sample to lysis tube, mix, heat, and incubate for seven minutes, (3) place on magnetic rack, allow beads to separate, aspirate supernatant, (4) add 100  $\mu$ l of Buffer 1, mix, place on magnetic rack, allow beads to separate, aspirate supernatant, (5) add 10  $\mu$ l of Buffer 2, mix, place in high temperature heat block for 3 minutes, place on magnetic rack, allow beads to separate, transfer 5  $\mu$ l supernatant, and (6) Add 5  $\mu$ l of Buffer 3, transfer 1 to 10  $\mu$ l of supernatant for PCR amplification and detection.

**[0192]** The PCR reagent mixture can be in the form of one or more lyophilized pellets and the steps by which the PCR-ready sample is prepared can involve reconstituting the PCR pellet by contacting it with liquid to create a PCR reagent mixture solution. In yet another embodiment, each of the PCR lanes may have dried down or lyophilized ASR reagents preloaded such that the user only needs to input prepared polynucleotide sample into the PCR. In another embodiment, the PCR lanes may have only the application-specific probes and primers pre-measured and pre-loaded, and the user inputs a sample mixed with the PCR reagents.

**[0193]** In various embodiments, the PCR-ready sample can include at least one probe that can be selective for a polynucleotide sequence, wherein the steps by which the PCR-ready sample is prepared involve contacting the neutralized polynucleotide sample or a PCR amplicon thereof with the probe. The probe can be a fluorogenic hybridization probe.

The fluorogenic hybridization probe can include a polynucleotide sequence coupled to a fluorescent reporter dye and a fluorescence quencher dye.

**[0194]** In various embodiments, the PCR-ready sample further includes a sample buffer.

**[0195]** In various embodiments, the PCR-ready sample includes at least one probe that is selective for a polynucleotide sequence, e.g., the polynucleotide sequence that is characteristic of a pathogen selected from the group consisting of gram positive bacteria, gram negative bacteria, yeast, fungi, protozoa, and viruses.

**[0196]** In various embodiments, the PCR reagent mixture can further include a polymerase enzyme, a positive control plasmid and a fluorogenic hybridization probe selective for at least a portion of the plasmid.

**[0197]** In various embodiments, the probe can be selective for a polynucleotide sequence that is characteristic of an organism, for example any organism that employs deoxyribonucleic acid or ribonucleic acid polynucleotides. Thus, the probe can be selective for any organism. Suitable organisms include mammals (including humans), birds, reptiles, amphibians, fish, domesticated animals, wild animals, extinct organisms, bacteria, fungi, viruses, plants, and the like. The probe can also be selective for components of organisms that employ their own polynucleotides, for example mitochondria. In some embodiments, the probe is selective for microorganisms, for example, organisms used in food production (for example, yeasts employed in fermented products, molds or bacteria employed in cheeses, and the like) or pathogens (e.g., of humans, domesticated or wild mammals, domesticated or wild birds, and the like). In some embodiments, the probe is selective for organisms selected from the group consisting of gram positive bacteria, gram negative bacteria, yeast, fungi, protozoa, and viruses.

**[0198]** In various embodiments, the probe can be selective for a polynucleotide sequence that is characteristic of an organism selected from the group consisting of *Staphylococcus* spp., e.g., *S. epidermidis*, *S. aureus*, Methicillin-resistant *Staphylococcus aureus* (MRSA), Vancomycin-resistant *Staphylococcus*; *Streptococcus* (e.g.,  $\alpha$ ,  $\beta$  or  $\gamma$ -hemolytic, Group A, B, C, D or G) such as *S. pyogenes*, *S. agalactiae*; *E. faecalis*, *E. durans*, and *E. faecium* (formerly *S. faecalis*, *S. durans*, *S. faecium*); nonenterococcal group D streptococci, e.g., *S. bovis* and *S. equines*; Streptococci viridans, e.g., *S. mutans*, *S. sanguis*, *S. salivarius*, *S. mitior*, *A. milleri*, *S. constellatus*, *S. intermedius*, and *S. anginosus*; *S. intiae*; *S. pneumoniae*; *Neisseria*, e.g., *N. meningitidis*, *N. gonorrhoeae*, saprophytic *Neisseria* sp; *Erysipelothrix*, e.g., *E. rhusiopathiae*; *Listeria* spp., e.g., *L. monocytogenes*, rarely *L. ivanovii* and *L. seeligeri*; *Bacillus*, e.g., *B. anthracis*, *B. cereus*, *B. subtilis*, *B. subtilis niger*, *B. thuringiensis*; *Nocardia* asteroides; *Legionella*, e.g., *L. pneumophila*, *Pneumocystis*, e.g., *P. carinii*; Enterobacteriaceae such as *Salmonella*, *Shigella*, *Escherichia* (e.g., *E. coli*, *E. coli*O157:H7); *Klebsiella*, *Enterobacter*, *Serratia*, *Proteus*, *Morganella*, *Providencia*, *Yersinia*, and the like, e.g., *Salmonella*, e.g., *S. typhi* *S. paratyphi* A, B (*S. schottmuelleri*), and C (*S. hirschfeldii*), *S. dublin*, *S. choleraesuis*, *S. enteritidis*, *S. typhimurium*, *S. heidelberg*, *S. newport*, *S. infantis*, *S. agona*, *S. montevideo*, and *S. saint-paul*; *Shigella* e.g., subgroups: A, B, C, and D, such as *S. flexneri*, *S. sonnei*, *S. boydii*, *S. dysenteriae*; *Proteus* (*P. mirabilis*, *P. vulgaris*, and *P. myxofaciens*), *Morganella* (*M. morganii*); *Providencia* (*P. rettgeri*, *P. alcalifaciens*, and *P. stuartii*); *Yersinia*, e.g., *Y. pestis*, *Y.*