

[0152] Any of the isolated nucleic acid molecules of the present invention may be introduced into a plant cell in a permanent or transient manner in combination with other genetic elements such as vectors, promoters, enhancers etc. Further any of the nucleic acid molecules encoding a *D. v. virgifera* protein or fragment thereof may be introduced into a plant cell in a manner that allows for over expression of the protein or polypeptide encoded by the nucleic acid molecule.

[0153] Antibodies have been expressed in plants (Hiatt et al., *Nature* 342:76-78 (1989); Conrad and Fielder, *Plant Mol. Biol.* 26:1023-1030 (1994)). Cytoplasmic expression of a scFv (single-chain Fv antibodies) has been reported to delay infection by artichoke mottled crinkle virus. Transgenic plants that express antibodies directed against endogenous proteins may exhibit a physiological effect (Philips et al., *EMBO J.* 16:4489-4496 (1997); Marion-Poll, *Trends in Plant Science* 2:447-448 (1997)). For example, expressed anti-abscisic antibodies reportedly result in a general perturbation of seed development (Philips et al., *EMBO J.* 16:4489-4496 (1997)).

[0154] Antibodies that are catalytic may also be expressed in plants (abzymes). The principle behind abzymes is that since antibodies may be raised against many molecules, this recognition ability can be directed toward generating antibodies that bind transition states to force a chemical reaction forward (Persidas, *Nature Biotechnology* 15:1313-1315 (1997); Baca et al., *Ann. Rev. Biophys. Biomol. Struct.* 26:461-493 (1997)). The catalytic abilities of abzymes may be enhanced by site directed mutagenesis. Examples of abzymes are, for example, set forth in U.S. Pat. No. 5,658,753; U.S. Pat. No. 5,632,990; U.S. Pat. No. 5,631,137; U.S. Pat. No. 5,602,015; U.S. Pat. No. 5,559,538; U.S. Pat. No. 5,576,174; U.S. Pat. No. 5,500,358; U.S. Pat. No. 5,318,897; U.S. Pat. No. 5,298,409; U.S. Pat. No. 5,258,289 and U.S. Pat. No. 5,194,585, all of which are herein incorporated in their entirety.

[0155] It is understood that any of the antibodies of the present invention may be expressed in plants and that such expression can result in a physiological effect. It is also understood that any of the expressed antibodies may be catalytic.

[0156] (f) Microbial Constructs and Transformed Microbial Cells

[0157] The nucleotide sequences of the present invention may be introduced into a wide variety of prokaryotic and eukaryotic microorganism hosts to express the *D. v. virgifera* proteins or fragments thereof of interest. The term "microorganism" includes prokaryotic and eukaryotic microbial species such as bacteria and fungi. Fungi include yeast and filamentous fungi. Illustrative prokaryotes, both Gram-negative and Gram-positive, include Enterobacteriaceae, such as *Escherichia*, *Erwinia*, *Shigella*, *Salmonella*, and *Proteus*; Bacillaceae; Rhizobiceae, such as *Rhizobium*; Spirillaceae, such as *photobacterium*, *Zymomonas*, *Serratia*, *Aeromonas*, *Vibrio*, *Desulfovibrio*, *Spirillum*; Lactobacillaceae; Pseudomonadaceae, such as *Pseudomonas* and *Acetobacter*; Azotobacteraceae, Actinomycetales, and Nitrohacteraceae. Among eukaryotes are fungi, such as *Phycomycetes* and *Ascomycetes*, which includes yeast, such as *Saccharomyces* and *Schizosaccharomyces*; and *Basidiomycetes* yeast, such as *Rhodotorula*, *Aureobasidium*, *Sporobolomyces*, and the like.

[0158] For the purpose of plant protection against insects, a large number of microorganisms known to inhabit the phylloplane (the surface of the plant leaves) and/or the rhizosphere (the soil surrounding plant roots) of a wide variety of

important crops may also be desirable host cells for manipulation, propagation, storage, delivery and/or mutagenesis of the disclosed recombinant constructs. These microorganisms include bacteria, algae, and fungi. Of particular interest are microorganisms, such as bacteria, e.g., genera *Bacillus* (including the species and subspecies *B. thuringiensis kurstaki* HD-1, *B. thuringiensis kurstaki* HD-73, *B. thuringiensis sotto*, *B. thuringiensis berliner*, *B. thuringiensis thuringiensis*, *B. thuringiensis tolworthi*, *B. thuringiensis dendrolimus*, *B. thuringiensis alesti*, *B. thuringiensis galleriae*, *B. thuringiensis aizawai*, *B. thuringiensis subtoxicus*, *B. thuringiensis entomocidus*, *B. thuringiensis tenebrionis* and *B. thuringiensis san diego*); *Pseudomonas*, *Erwinia*, *Serratia*, *Klebsiella*, *Zanthomonas*, *Streptomyces*, *Rhizobium*, *Rhodopseudomonas*, *Methylophilus*, *Agrobacterium*, *Acetobacter*, *Lactobacillus*, *Arthrobacter*, *Azotobacter*, *Leuconostoc*, and *Alcaligenes*; fungi, particularly yeast, e.g., genera *Saccharomyces*, *Cryptococcus*, *Kluyveromyces*, *Sporobolomyces*, *Rhodotorula*, and *Aureobasidium*. Of particular interest are such phytosphere bacterial species as *Pseudomonas syringae*, *Pseudomonas fluorescens*, *Serratia marcescens*, *Acetobacter xylinum*, *Agrobacterium tumefaciens*, *Rhodobacter sphaeroides*, *Xanthomonas campestris*, *Rhizobium melioli*, *Alcaligenes eutrophus*, and *Azotobacter vinlandii*; and phytosphere yeast species such as *Rhodotorula rubra*, *R. glutinis*, *R. marina*, *R. aurantiaca*, *Cryptococcus albidus*, *C. diffluens*, *C. laurentii*, *Saccharomyces rosei*, *S. pretoriensis*, *S. cerevisiae*, *Sporobolomyces roseus*, *S. odoratus*, *Kluyveromyces veronae*, and *Aureobasidium pollulans*.

[0159] It is well known that exogenous nucleic acids encoding polypeptides of interest can be introduced into a microbial host cell, such as a bacterial cell or a fungal cell, using a recombinant construct. The present invention also relates to a fungal or bacterial recombinant construct comprising a structural nucleotide sequence encoding a *D. v. virgifera* protein or fragment thereof. The present invention also relates to a bacterial or fungal cell comprising a bacterial or fungal recombinant vector of the present invention. The present invention also relates to methods for obtaining a recombinant bacterial or fungal host cell, comprising introducing into a bacterial or fungal host cell an exogenous nucleic acid molecule of the present invention.

[0160] The bacterial recombinant vector may be a linear or a closed circular plasmid. The vector system may be a single vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the bacterial host. In addition, the bacterial vector may be an expression vector. Nucleic acid molecules encoding *D. v. virgifera* proteins or fragments thereof can, for example, be suitably inserted into a replicable vector for expression in a bacterium under the control of a suitable promoter for that bacterium. Many vectors are available for this purpose, and selection of the appropriate vector will depend mainly on the size of the nucleic acid to be inserted into the vector and the particular host cell to be transformed with the vector. Each vector contains various components depending on its function (amplification of DNA or expression of DNA) and the particular host cell with which it is compatible. The vector components for bacterial transformation generally include, but are not limited to, one or more of the following: a signal sequence, an origin of replication, one or more selectable marker genes, and an inducible promoter allowing the expression of exogenous DNA.