


herein, or salt, solvate, or derivative thereof, (b) an antigen from a hemorrhagic fever virus; and (c) an adjuvant.

[0024] In some aspects, the hemorrhagic fever virus is a Filoviridae virus. In some aspects, the Filoviridae virus is Ebola virus. In another aspect, the adjuvant is an aluminum-containing adjuvant or MF59.

[0025] In another aspect of the invention, the benzonaphthyrindine TLR7 agonist is a benzonaphthyrindine of Formula (I) having the structure:



Formula (I)

wherein:

[0026] R^3 is H, halogen, C_1 - C_6 alkyl, C_2 - C_8 alkene, C_2 - C_8 alkyne, C_1 - C_6 heteroalkyl, C_1 - C_6 haloalkyl, C_1 - C_6 alkoxy, C_1 - C_6 haloalkoxy, aryl, heteroaryl, C_3 - C_8 cycloalkyl, and C_3 - C_8 heterocycloalkyl, wherein the C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 haloalkyl, C_1 - C_6 alkoxy, C_1 - C_6 haloalkoxy, C_3 - C_8 cycloalkyl, or C_3 - C_8 heterocycloalkyl groups of R^3 are each optionally substituted with 1 to 3 substituents independently selected from halogen, $-CN$, $-R^7$, $-OR^8$, $-C(O)R^8$, $-OC(O)R^8$, $-C(O)OR^8$, $-N(R^9)_2$, $-C(O)N(R^9)_2$, $-S(O)_2R^8$, $-S(O)_2N(R^9)_2$ and $-NR^9S(O)_2R^8$;

[0027] R^4 and R^5 are each independently selected from H, halogen, $-C(O)OR^7$, $-C(O)R^7$, $-C(O)N(R^{11}R^{12})$, $-N(R^{11}R^{12})$, $-N(R^9)_2$, $-NHN(R^9)_2$, $-SR^7$, $-(CH_2)_m$, $-OR^7$, $-(CH_2)_mR^7$, $-LR^8$, $-LR^{10}$, $-OLR^8$, $-OLR^{10}$, C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 haloalkyl, C_2 - C_8 alkene, C_2 - C_8 alkyne, C_1 - C_6 alkoxy, C_1 - C_6 haloalkoxy, aryl, heteroaryl, C_3 - C_8 cycloalkyl, and C_3 - C_8 heterocycloalkyl, wherein the C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 haloalkyl, C_2 - C_8 alkene, C_2 - C_8 alkyne, C_1 - C_6 alkoxy, C_1 - C_6 haloalkoxy, aryl, heteroaryl, C_3 - C_8 cycloalkyl, and C_3 - C_8 heterocycloalkyl groups of R^4 and R^5 are each optionally substituted with 1 to 3 substituents independently selected from halogen, $-CN$, $-NO_2$, $-R^7$, $-OR^8$, $-C(O)R^8$, $-OC(O)R^8$, $-C(O)OR^8$, $-N(R^9)_2$, $-P(O)(OR^8)_2$, $-OP(O)(OR^8)_2$, $-P(O)(OR^{10})_2$, $-OP(O)(OR^{10})_2$, $-C(O)N(R^9)_2$, $-S(O)_2R^8$, $-S(O)R^8$, $-S(O)_2N(R^9)_2$, and $-NR^9S(O)_2R^8$;

[0028] or R^3 and R^4 , or R^4 and R^5 , when present on adjacent ring atoms, can optionally be linked together to form a 5-6 membered ring, wherein the 5-6 membered ring is optionally substituted with R^7 ;

[0029] each L is independently selected from a bond, $-(O(CH_2)_m)_t-$, C_1 - C_6 alkyl, C_2 - C_6 alkenylene and C_2 - C_6 alkynylene, wherein the C_1 - C_6 alkyl, C_2 - C_6 alkenylene and C_2 - C_6 alkynylene of L are each optionally substituted with 1 to 4 substituents independently selected from halogen, $-R^8$, $-OR^8$, $-N(R^9)_2$, $-P(O)(OR^8)_2$, $-OP(O)(OR^8)_2$, $-P(O)(OR^{10})_2$, and $-OP(O)(OR^{10})_2$;

[0030] R^7 is selected from H, C_1 - C_6 alkyl, aryl, heteroaryl, C_3 - C_8 cycloalkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 haloalkyl,

C_2 - C_8 alkene, C_2 - C_8 alkyne, C_1 - C_6 alkoxy, C_1 - C_6 haloalkoxy, and C_3 - C_8 heterocycloalkyl, wherein the C_1 - C_6 alkyl, aryl, heteroaryl, C_3 - C_8 cycloalkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 haloalkyl, C_2 - C_8 alkene, C_2 - C_8 alkyne, C_1 - C_6 alkoxy, C_1 - C_6 haloalkoxy, and C_3 - C_8 heterocycloalkyl groups of R^7 are each optionally substituted with 1 to 3 R^{13} groups;

[0031] each R^8 is independently selected from H, $-CH(R^{10})_2$, C_1 - C_8 alkyl, C_2 - C_8 alkene, C_2 - C_8 alkyne, C_1 - C_6 haloalkyl, C_1 - C_6 alkoxy, C_1 - C_6 heteroalkyl, C_3 - C_8 cycloalkyl, C_2 - C_8 heterocycloalkyl, C_1 - C_6 hydroxyalkyl and C_1 - C_6 haloalkoxy, wherein the C_1 - C_8 alkyl, C_2 - C_8 alkene, C_2 - C_8 alkyne, C_1 - C_6 heteroalkyl, C_1 - C_6 haloalkyl, C_1 - C_6 alkoxy, C_3 - C_8 cycloalkyl, C_2 - C_8 heterocycloalkyl, C_1 - C_6 hydroxyalkyl and C_1 - C_6 haloalkoxy groups of R^8 are each optionally substituted with 1 to 3 substituents independently selected from $-CN$, R^{11} , $-OR^{11}$, $-SR^{11}$, $-C(O)R^{11}$, $-OC(O)R^{11}$, $-C(O)N(R^9)_2$, $-C(O)OR^{11}$, $-NR^9C(O)R^{11}$, $-NR^9R^{10}$, $-NR^{11}R^{12}$, $-N(R^9)_2$, $-OR^9$, $-OR^{10}$, $-C(O)NR^{11}R^{12}$, $-C(O)NR^{11}OH$, $-S(O)_2R^{11}$, $-S(O)R^{11}$, $-S(O)_2NR^{11}R^{12}$, $-NR^{11}S(O)_2R^{11}$, $-P(O)(OR^{11})_2$, and $-OP(O)(OR^{11})_2$;

[0032] each R^9 is independently selected from H, $-C(O)R^8$, $-C(O)OR^8$, $-C(O)R^{10}$, $-C(O)OR^{10}$, $-S(O)_2R^{10}$, $-C_1$ - C_6 alkyl, C_1 - C_6 heteroalkyl and C_3 - C_6 cycloalkyl, or each R^9 is independently a C_1 - C_6 alkyl that together with N they are attached to form a C_3 - C_8 heterocycloalkyl, wherein the C_3 - C_8 heterocycloalkyl ring optionally contains an additional heteroatom selected from N, O and S, and wherein the C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_3 - C_6 cycloalkyl, or C_3 - C_8 heterocycloalkyl groups of R^9 are each optionally substituted with 1 to 3 substituents independently selected from $-CN$, R^{11} , $-OR^{11}$, $-SR^{11}$, $-C(O)R^{11}$, $-OC(O)R^{11}$, $-C(O)OR^{11}$, $-NR^{11}R^{12}$, $-C(O)NR^{11}R^{12}$, $-C(O)NR^{11}OH$, $-S(O)_2R^{11}$, $-S(O)R^{11}$, $-S(O)_2NR^{11}R^{12}$, $-NR^{11}S(O)_2R^{11}$, $-P(O)(OR^{11})_2$, and $-OP(O)(OR^{11})_2$;

[0033] each R^{10} is independently selected from aryl, C_3 - C_8 cycloalkyl, C_3 - C_8 heterocycloalkyl and heteroaryl, wherein the aryl, C_3 - C_8 cycloalkyl, C_3 - C_8 heterocycloalkyl and heteroaryl groups are optionally substituted with 1 to 3 substituents selected from halogen, $-R^8$, $-OR^8$, $-LR^9$, $-LOR^9$, $-N(R^9)_2$, $-NR^9C(O)R^8$, $-NR^9CO_2R^8$, $-CO_2R^8$, $-C(O)R^8$ and $-C(O)N(R^9)_2$;

[0034] R^{11} and R^{12} are independently selected from H, C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 haloalkyl, aryl, heteroaryl, C_3 - C_8 cycloalkyl, and C_3 - C_8 heterocycloalkyl, wherein the C_1 - C_6 alkyl, C_1 - C_6 heteroalkyl, C_1 - C_6 haloalkyl, aryl, heteroaryl, C_3 - C_8 cycloalkyl, and C_3 - C_8 heterocycloalkyl groups of R^{11} and R^{12} are each optionally substituted with 1 to 3 substituents independently selected from halogen, $-CN$, R^8 , $-OR^8$, $-C(O)R^8$, $-OC(O)R^8$, $-C(O)OR^8$, $-N(R^9)_2$, $-NR^8C(O)R^8$, $-NR^8C(O)OR^8$, $-C(O)N(R^9)_2$, C_3 - C_8 heterocycloalkyl, **[0035]** $-S(O)_2R^8$, $-S(O)_2N(R^9)_2$, $-NR^8S(O)_2R^8$, C_1 - C_6 haloalkyl and C_1 - C_6 haloalkoxy;

[0036] or R^{11} and R^{12} are each independently C_1 - C_6 alkyl and taken together with the N atom to which they are attached form an optionally substituted C_3 - C_8 heterocycloalkyl ring optionally containing an additional heteroatom selected from N, O and S;

[0037] each R^{13} is independently selected from halogen, $-CN$, $-LR^9$, $-LOR^9$, $-OLR^9$, $-LR^{10}$, $-LOR^{10}$, $-OLR^{10}$,