

**[0016]** Definitions

**[0017]** “Small molecule”, as used herein, means a molecule less than 5 kiloDalton, more typically less than 1 kiloDalton. As used herein, “small molecule” excludes proteins.

**[0018]** The term “candidate drug” as used herein, refers to any medicinal substance used in humans, animals, or plants. Encompassed within this definition are compound analogs, naturally occurring, synthetic and recombinant pharmaceuticals, hormones, antimicrobials, neurotransmitters, etc. This includes any substance or precursor (whether naturally occurring, synthetic or recombinant) which is to be evaluated for use as a drug for treatment of neurodegenerative disease, or other disease characterized by aberrant aggregation, or prevention thereof. Evaluation typically takes place through activity in an assay, such as the screening assays of the present invention.

**[0019]** A variety of types of particles can be used in the invention. For example, “fluid suspendable particle” means a particle that can be made to stay in suspension in a fluid in which it is used for purposes of the invention (typically an aqueous solution) by itself, or can be maintained in solution by application of a magnetic field, an electromagnetic field, agitation such as stirring, shaking, vibrating, sonicating, centrifuging, vortexing, or the like. A “magnetically suspendable” particle is one that can be maintained in suspension in a fluid via application of a magnetic field. An electromagnetically-suspendable particle is one that can be maintained in suspension in a fluid by application of an electromagnetic field (e.g., a particle carrying a charge, or a particle modified to carry a charge). A “self-suspendable particle” is a particle that is of low enough size and/or mass that it will remain in suspension in a fluid in which it is used (typically an aqueous solution), without assistance of, for example, a magnetic field, for at least 1 hour. Other self-suspendable particles will remain in suspension, without assistance, for 5 hours, 1 day, 1 week, or even 1 month, in accordance with the invention.

**[0020]** “Proteins” and “peptides” are well-known terms in the art, and are not precisely defined in the art in terms of the number of amino acids that each includes. As used herein, these terms are given their ordinary meaning in the art. Generally, peptides are amino acid sequences of less than about 100 amino acids in length, but can include sequences of up to 300 amino acids. Proteins generally are considered to be molecules of at least 100 amino acids.

**[0021]** As used herein, a “metal binding tag” refers to a group of molecules that can become fastened to a metal that is coordinated by a chelate. Suitable groups of such molecules include amino acid sequences, typically from about 2 to about 10 amino acid residues. These include, but are not limited to, histidines and cysteines (“polyamino acid tags”). Such binding tags, when they include histidine, can be referred to as a “polyhistidine tract” or “histidine tag” or “HIS-tag”, and can be present at either the amino- or carboxy-terminus, or at any exposed region, of a peptide or protein or nucleic acid. A poly-histidine tract of six to ten residues is preferred for use in the invention. The polyhistidine tract is also defined functionally as being a number of consecutive histidine residues added to a protein of interest which allows the affinity purification of the resulting protein on a metal chelate column, or the identification of a protein

terminus through the interaction with another molecule (e.g. an antibody reactive with the HIS-tag).

**[0022]** “Affinity tag” is given its ordinary meaning in the art. Affinity tags include, for example, metal binding tags, GST (in GST/glutathione binding clip), and streptavidin (in biotin/streptavidin binding). At various locations herein specific affinity tags are described in connection with binding interactions. It is to be understood that the invention involves, in any embodiment employing an affinity tag, a series of individual embodiments each involving selection of any of the affinity tags described herein.

**[0023]** As used herein, “chelate coordinating a metal” or metal coordinated by a chelate, refers to a metal coordinated by a chelating agent that does not fill all available coordination sites on the metal, leaving some coordination sites available for binding via a metal binding tag.

**[0024]** As used herein, “metal binding tag/metal/chelate linkage” defines a linkage between first and second species in which a first species is immobilized relative to a metal binding tag and a second species is immobilized relative to a chelate, where the chelate coordinates a metal to which the metal binding tag is also coordinated. U.S. Pat. No. 5,620,850 of Bamdad, et al., incorporated herein by reference, describes exemplary linkages.

**[0025]** “Signaling entity” means an entity that is capable of indicating its existence in a particular sample or at a particular location. Signaling entities of the invention can be those that are identifiable by the unaided human eye, those that may be invisible in isolation but may be detectable by the unaided human eye if in sufficient quantity (e.g., colloid particles), entities that absorb or emit electromagnetic radiation at a level or within a wavelength range such that they can be readily detected visibly (unaided or with a microscope including an electron microscope or the like), or spectroscopically, entities that can be detected electronically or electrochemically, such as redox-active molecules exhibiting a characteristic oxidation/reduction pattern upon exposure to appropriate activation energy (“electronic signaling entities”), or the like. Examples include dyes, pigments, electroactive molecules such as redox-active molecules, fluorescent moieties (including, by definition, phosphorescent moieties), up-regulating phosphors, chemiluminescent entities, electrochemiluminescent entities, or enzyme-linked signaling moieties including horse radish peroxidase and alkaline phosphatase. “Precursors of signaling entities” are entities that by themselves may not have signaling capability but, upon chemical, electrochemical, electrical, magnetic, or physical interaction with another species, become signaling entities. An example includes a chromophore having the ability to emit radiation within a particular, detectable wavelength only upon chemical interaction with another molecule. Precursors of signaling entities are distinguishable from, but are included within the definition of, “signaling entities” as used herein. As used herein, “fastened to or adapted to be fastened”, in the context of a species relative to another species or to a surface of an article, means that the species is chemically or biochemically linked via covalent attachment, attachment via specific biological binding (e.g., biotin/streptavidin), coordinative bonding such as chelate/metal binding, or the like. For example, “fastened” in this context includes multiple chemical linkages, multiple chemical/biological linkages, etc., including, but not limited