

saya, cauliflower, cedar, a cereal, celery, chestnut, cherry, chinese cabbage, citrus, clemantine, clover, coffee, corn, cotton, cowpea, cucumber, cypress, eggplant, elm, endive, eucalyptus, fennel, figs, fir, geranium, grape, grapefruit, groundnuts, ground cherry, gum hemlock, hickory, kale, kiwifruit, kohlrabi, larch, lettuce, leek, lemon, lime, locust, pine, maidenhair, maize, mango, maple, melon, millet, mushroom, mustard, nuts, oak, oats, okra, onion, orange, an ornamental plant or flower or tree, papaya, palm, parsley, parsnip, pea, peach, peanut, pear, peat, pepper, persimmon, pigeon pea, pine, pineapple, plantain, plum, pomegranate, potato, pumpkin, radicchio, radish, rapeseed, raspberry, rice, rye, sorghum, soybean, spinach, spruce, squash, strawberry, sugarbeet, sugarcane, sunflower, sweet potato, sweet corn, tangerine, tea, tobacco, tomato, trees, triticale, turf grasses, turnips, a vine, walnut, watercress, watermelon, wheat, yams, yew, and zucchini.

**[0081]** According to the present invention “plant tissue” also encompasses plant cells. Plant cells include suspension cultures, callus, embryos, meristematic regions, callus tissue, leaves, roots, shoots, gametophytes, sporophytes, pollen, seeds and microspores. Plant tissues may be at various stages of maturity and may be grown in liquid or solid culture, or in soil or suitable media in pots, greenhouses or fields. A plant tissue also refers to any clone of such a plant, seed, progeny, propagule whether generated sexually or asexually, and descendants of any of these, such as cuttings or seed.

**[0082]** Promoter is intended to mean a nucleic acid, preferably DNA that binds RNA polymerase and/or other transcription regulatory elements. As with any promoter, the promoters of the current invention will facilitate or control the transcription of DNA or RNA to generate an mRNA molecule from a nucleic acid molecule that is operably linked to the promoter. As stated earlier, the RNA generated may code for a protein or polypeptide or may code for an RNA interfering, or antisense molecule.

**[0083]** Polynucleotide is a nucleotide sequence, comprising a gene coding sequence or a fragment thereof, a promoter, an intron, an enhancer region, a polyadenylation site, a translation initiation site, 5' or 3' untranslated regions, a reporter gene, a selectable marker or the like. The polynucleotide may comprise single stranded or double stranded DNA or RNA. The polynucleotide may comprise modified bases or a modified backbone. The polynucleotide may be genomic, an RNA transcript (such as an mRNA) or a processed nucleotide sequence (such as a cDNA). The polynucleotide may comprise a sequence in either sense or antisense orientations.

**[0084]** An isolated polynucleotide is a polynucleotide sequence that is not in its native state, e.g., the polynucleotide is comprised of a nucleotide sequence not found in nature or the polynucleotide is separated from nucleotide sequences with which it typically is in proximity or is next to nucleotide sequences with which it typically is not in proximity.

**[0085]** Recombinant nucleotide sequence refers to a nucleic acid molecule that contains a genetically engineered modification through manipulation via mutagenesis, restriction enzymes, and the like.

**[0086]** RNA interference (RNAi) refers to sequence-specific or gene-specific suppression of gene expression (protein synthesis) that is mediated by short interfering RNA (siRNA).

**[0087]** Sequence identity: as used herein, “sequence identity” or “identity” in the context of two nucleic acid sequences

includes reference to the residues in the two sequences which are the same when aligned for maximum correspondence over a specified region.

**[0088]** As used herein, percentage of sequence identity means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

**[0089]** “Sequence identity” has an art-recognized meaning and can be calculated using published techniques. See COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, ed. (Oxford University Press, 1988), BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, ed. (Academic Press, 1993), COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin & Griffin, eds., (Humana Press, 1994), SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, Von Heinje ed., Academic Press (1987), SEQUENCE ANALYSIS PRIMER, Gribkov & Devereux, eds. (Macmillan Stockton Press, 1991), and Carillo & Lipton, *SIAM J. Applied Math.* 48: 1073 (1988). Methods commonly employed to determine identity or similarity between two sequences include but are not limited to those disclosed in GUIDE TO HUGE COMPUTERS, Bishop, ed., (Academic Press, 1994) and Carillo & Lipton, supra. Methods to determine identity and similarity are codified in computer programs. Preferred computer program methods to determine identity and similarity between two sequences include but are not limited to the GCG program package (Devereux et al., *Nucleic Acids Research* 12: 387 (1984)), BLASTP, BLASTN, FASTA (Atschul et al., *J. Mol. Biol.* 215: 403 (1990)), and FASTDB (Brutlag et al., *Comp. App. Biosci.* 6: 237 (1990)).

**[0090]** Short hairpin RNA (shRNA) are short single-stranded RNAs having a high degree of secondary structure such that a portion of the RNA strand forms a hairpin loop.

**[0091]** Short interfering RNA (siRNA) refers to double-stranded RNA molecules from about 10 to about 30 nucleotides long that are named for their ability to specifically interfere with gene protein expression.

**[0092]** Target sequence refers to a nucleotide sequence in a pest that is selected for suppression or inhibition by double stranded RNA technology. A target sequence encodes an essential feature or biological activity within a pest.

**[0093]** Transcriptional terminators: The expression DNA constructs of the present invention typically have a transcriptional termination region at the opposite end from the transcription initiation regulatory region. The transcriptional termination region may be selected, for stability of the mRNA to enhance expression and/or for the addition of polyadenylation tails added to the gene transcription product. Translation of a nascent polypeptide undergoes termination when any of the three chain-termination codons enters the A site on the ribosome. Translation termination codons are UAA, UAG, and UGA.

**[0094]** Transformation: A process by which a nucleic acid is stably inserted into the genome of an organism. Transformation may occur under natural or artificial conditions using