

in significantly higher BPH mortalities at final concentrations of as low as 0.04 mg dsRNA per ml diet when compared with survival on diet only, as shown in FIG. 2-NL and Table 9-NL. Table 9-NL summarizes the survival of *Nilaparvata lugens* artificial diet feeding trial supplemented with 1, 0.2, 0.08, & 0.04 mg/ml (final concentration) of target NL002. In the survival analysis column the effect of RNAi is indicated as follows: +=significantly decreases survival compared to diet only control (alpha <0.05); -=no significant differences in survival compared to diet only control. Survival curves were compared using the logrank test.

F. Cloning of a BPH Gene Fragment in a Vector Suitable for Bacterial Production of Insect-Active Double-Stranded RNA

[0400] What follows is an example of cloning a DNA fragment corresponding to a BPH gene target in a vector for the expression of double-stranded RNA in a bacterial host, although any vector comprising a T7 promoter or any other promoter for efficient transcription in bacteria, may be used (reference to WO0001846).

[0401] The sequences of the specific primers used for the amplification of target genes are provided in Table 8. The template used is the pCR8/GW/topo vector containing any of target sequences. The primers are used in a PCR reaction with the following conditions: 5 minutes at 98° C., followed by 30 cycles of 10 seconds at 98° C., 30 seconds at 55° C. and 2 minutes at 72° C., followed by 10 minutes at 72° C. The resulting PCR fragment is analyzed on agarose gel, purified (QIAquick Gel Extraction kit, Cat. Nr. 28706, Qiagen), blunt-ended cloned into Srf I-linearized pGNA49A vector (reference to WO00188121A1), and sequenced. The sequence of the resulting PCR product corresponds to the respective sequence as given in Table 8-NL. The recombinant vector harbouring this sequence is named pGBNJ00.

G. Expression and Production of a Double-Stranded RNA Target in Two Strains of *Escherichia coli*: (1) AB309-105, and, (2) BL21(DE3)

[0402] The procedures described below are followed in order to express suitable levels of insect-active double-stranded RNA of insect target in bacteria. An RNaseIII-deficient strain, AB309-105, is used in comparison to wild-type RNaseIII-containing bacteria, BL21(DE3).

Transformation of AB309-105 and BL21(DE3)

[0403] Three hundred ng of the plasmid are added to and gently mixed in a 50 µl aliquot of ice-chilled chemically competent *E. coli* strain AB309-105 or BL21(DE3). The cells are incubated on ice for 20 minutes before subjecting them to a heat shock treatment of 37° C. for 5 minutes, after which the cells are placed back on ice for a further 5 minutes. Four hundred and fifty µl of room temperature SOC medium is added to the cells and the suspension incubated on a shaker (250 rpm) at 37° C. for 1 hour. One hundred µl of the bacterial cell suspension is transferred to a 500 ml conical flask containing 150 ml of liquid Luria-Bertani (LB) broth supplemented with 100 µg/ml carbenicillin antibiotic. The culture is incubated on an Innova 4430 shaker (250 rpm) at 37° C. overnight (16 to 18 hours).

Chemical Induction of Double-Stranded RNA Expression in AB309-105 and BL21(DE3)

[0404] Expression of double-stranded RNA from the recombinant vector, pGBNJ003, in the bacterial strain AB309-105 or BL21(DE3) is made possible since all the genetic components for controlled expression are present. In the presence of the chemical inducer isopropylthiogalactoside, or IPTG, the T7 polymerase will drive the transcription of the target sequence in both antisense and sense directions since these are flanked by oppositely oriented T7 promoters. **[0405]** The optical density at 600 nm of the overnight bacterial culture is measured using an appropriate spectrophotometer and adjusted to a value of 1 by the addition of fresh LB broth. Fifty ml of this culture is transferred to a 50 ml Falcon tube and the culture then centrifuged at 3000 g at 15° C. for 10 minutes. The supernatant is removed and the bacterial pellet resuspended in 50 ml of fresh S complete medium (SNC medium plus 5 µg/ml cholesterol) supplemented with 100 µg/ml carbenicillin and 1 mM IPTG. The bacteria are induced for 2 to 4 hours at room temperature.

Heat Treatment of Bacteria

[0406] Bacteria are killed by heat treatment in order to minimise the risk of contamination of the artificial diet in the test plates. However, heat treatment of bacteria expressing double-stranded RNA is not a prerequisite for inducing toxicity towards the insects due to RNA interference. The induced bacterial culture is centrifuged at 3000 g at room temperature for 10 minutes, the supernatant discarded and the pellet subjected to 80° C. for 20 minutes in a water bath. After heat treatment, the bacterial pellet is resuspended in 1.5 ml MilliQ water and the suspension transferred to a microfuge tube. Several tubes are prepared and used in the bioassays for each refreshment. The tubes are stored at -20° C. until further use.

H. Laboratory Trials to Test *Escherichia coli* Expressing dsRNA Targets Against *Nilaparvata lugens*

Plant-Based Bioassays

[0407] Whole plants are sprayed with suspensions of chemically induced bacteria expressing dsRNA prior to feeding the plants to BPH. The are grown from in a plant growth room chamber. The plants are caged by placing a 500 ml plastic bottle upside down over the plant with the neck of the bottle firmly placed in the soil in a pot and the base cut open and covered with a fine nylon mesh to permit aeration, reduce condensation inside and prevent insect escape. BPH are placed on each treated plant in the cage. Plants are treated with a suspension of *E. coli* AB309-105 harbouring the pGBNJ001 plasmids or pGN29 plasmid. Different quantities of bacteria are applied to the plants: for instance 66, 22, and 7 units, where one unit is defined as 10⁹ bacterial cells in 1 ml of a bacterial suspension at optical density value of 1 at 600 nm wavelength. In each case, a total volume of between 1 and 10 ml s sprayed on the plant with the aid of a vaporizer. One plant is used per treatment in this trial. The number of survivors are counted and the weight of each survivor recorded.

[0408] Spraying plants with a suspension of *E. coli* bacterial strain AB309-105 expressing target dsRNA from pGBNJ003 led to a dramatic increase in insect mortality when compared to pGN29 control. These experiments show