before thyroidectomy: Final results, Ann Surg. 2008 May; 247(5):843-53. [9] Stojadinovic A, Fields S I, Shriver C D, et al., Electrical Impedance Scanning of Thyroid Nodules Prior to Thyroid Surgery: A Prospective Study, Annals of Surgical Oncology 2005; 12(2): 152-160. [10] Raza S N, Shah M D, Palme C E, et al., Risk factors for well-differentiated thyroid carcinoma in patients with thyroid nodular disease, Otolaryngol Head Neck Surg. 2008 July; 139(1):21-6. [11] Varverakis E, Neonakis E, Tzardi M, Chrysos E., Role of color Doppler ultrasonography in the preoperative management of cold thyroid nodules, Hormones 2007; 6(1):44-51. [12] Lyshchik A, Moses R, Barnes S L, et al., Quantitative analysis of tumor vascularity in benign and malignant solid thyroid nodules, J Ultrasound Med. 2007 June; 26(6):837-46. [13] Varverakis E, Neonakis E., Contribution of high-resolution ultrasonography in the differential diagnosis of benign from malignant thyroid nodules, Hormones (Athens) 2002; 1: 51-56. [14] Spiezia S, Cerbone G, Colao A, et al., Usefulness of power Doppler in the diagnostic management of hypoechoic thyroid nodules, Eur J Ultrasound 1997; 6: 165-170. [15] Rago T, Vitti P, Chiovato L et al., Role of conventional ultrasonography and color flowdoppler sonography in predicting malignancy in 'cold' thyroid nodules, Eur J Endocrinol 1998 January; 138(1):41-6. [16] Bae U, Dighe M, Dubinsky T, et al., Ultrasound thyroid elastography using carotid artery pulsation: preliminary study, J Ultrasound Med. 2007 June; 26(6): 797-805. [17] Rago T, Santini F, Scutari M, Pinchera A, Vitti P., Elastography: new developments in ultrasound for predicting malignancy in thyroid nodules, J Clin Endocrinol Metab. 2007 August; 92(8):2917-22. [18] Sebastianes F M, Cerci J J, Zanoni P H, et al., Role of 18F-FDG PET in Preoperative Assessment of Cytologically Indeterminate Thyroid Nodules, J Clin Endocrinol Metab. 2007 August 7. [19] De Geus-Oei L F, Pieters G F F M, Bonenkamp J J, et al, 18 F-FDG PET reduces unnecessary hemithyroidectomies for thyroid nodules with indeterminate cytologic results, J Nucl Med 2006; 47:770-775. [20] Fricke H, Morse S., The electric capacity of tumors of the breast, J Cancer Res 16:310-376. 1926. [21] Scholz B, Anderson R., On electrical impedance scanning—principles and simulations, Electromedica 2000; 68:35-44. [22] Glickman Y A, Filo O, David M, Yayon A, Topaz M, Zamir B, Ginzburg A, Rozenman D, Kenan G., Electrical impedance scanning: a new approach to skin cancer diagnosis, Skin Res Technol. 2003 August; 9(3):262-8. [23] Malich A, Fritsch T, Mauch C, et al., Electrical impedance scanning: a new technique in the diagnosis of lymph nodes in which malignancy is suspected on ultrasound, Br J Radiol. 2001; 74(877):42-7. [24] Malich A, Boehm T, Facius M, et al., Use of electrical impedance scanning in the differentiation of sonographically suspicious and highly suspicious lymph nodes of the headneck region, Eur Radiol. 2002; 12(5):1114-20. [25] Mentzel H J, Malich A, Kentouche K, Freesmeyer M, Bottcher J, Schneider G, Gruhn B, Vogt S, Zintl F, Anderson R, Kaiser W A., Electrical impedance scanning-application of this new technique for lymph node evaluation in children, Pediatr Radiol. 2003; 33(7):461-6. [26] Malich A, Fritsch T, Anderson R, et al., Electrical impedance scanning for classifying suspicious breast lesions: first results, Eur Radiol. 2000; 10(10):1555-61. [27] Malich A, Boehm T, Facius M., Differentation of mammographically suspicious lesions: evaluation of breast ultrasound, MRI mammography and electrical impedance scanning as adjunctive technologies in breast cancer detection, Clinical Radiology 2001; 56: 278-83. [28] Fuchsjaeger M H, Flory D, Reiner C S, Rudas M, Riedl C C, Helbich T H, The negative predictive value of electrical impedance scanning in BI-RADS category IV breast lesions, Invest Radiol. 2005; 40(7):478-85. [29] Stojadinovic A, Nissan A, Gallimidi Z, et al., Electrical Impedance Scanning for the Early Detection of Breast Cancer in Young Women: Preliminary Results of a Multi-Center Prospective Clinical Trial, J Clin Oncol 2005; 23(12):2703-2715. [30] Stojadinovic A, Moskovitz O, Gallimidi Z, et al., Prospective Study of Electrical Impedance Scanning for Identifying Young Women at Risk for Breast Cancer, Br Cancer Res Treat 2006; 97(2): 179-89. [31] Tuttle R M, Lemar H, Burch H B, Clinical features associated with an increased risk of thyroid malignancy in patients with follicular neoplasia by fine-needle aspiration, Thyroid 1998 May; 8(5):377-83. [32] Montgomery, S. P., Swiecki, C. W. & Shriver, C. D., The evaluation of casualties from Operation Iraqi Freedom on return to the continental United States from March to June 2003, *J Am Coll Surg* 201:7-12; discussion 12-13 (2005). [33] Peoples, G. E., Jezior, J. R. & Shriver, C. D., Caring for the wounded in Iraq—a photo essay, *N Engl J Med* 351:2476-2480 (2004). [34] Owens, B. D., et al., Combat wounds in operation Iraqi Freedom and operation Enduring Freedom, *J Trauma* 64:295-299 (2008). [35] Marsh, D. J., Abu-Sitta, G. & Patel, H., The role of vacuum-assisted wound closure in blast injury, *Plast Reconstr Surg* 119:1978-1979 (2007). [36] Breugem, C. C. & Strackee, S. D., Is there evidence-based guidance for timing of soft tissue coverage of grade III B tibia fractures? *Int J Low Extrem Wounds* 5:261-270 (2006). [37] Nwomeh, B. C., Yager, D. R. & Cohen, I. K., Physiology of the chronic wound, *Clin Plast Surg* 25:341-356 (1998). [38] DeLong, W. G., Jr. & Born, C. T., Cytokines in patients with polytrauma, *Clin Orthop Relat Res:* 57-65 (2004). [39] Nast-Kolb, D., et al., Indicators of the posttraumatic inflammatory response correlate with organ failure in patients with multiple injuries, *J Trauma* 42, 446-454; discussion 454-445 (1997). [40] Endo, S., et al., Plasma endotoxin and cytokine concentrations in patients with hemorrhagic shock, *Crit Care Med* 22, 949-955 (1994). [41] Martin, C., Boisson, C., Haccoun, M., Thomachot, L. & Mege, J. L., Patterns of cytokine evolution (tumor necrosis factor-alpha and interleukin-6) after septic shock, hemorrhagic shock, and severe trauma, *Crit Care Med* 25, 1813-1819 (1997). [42] Biffl, W. L., Moore, E. E., Moore, F. A. & Peterson, V. M., Interleukin-6 in the injured patient. Marker of injury or mediator of inflammation? *Ann Surg* 224, 647-664 (1996). [43] Neidhardt, R., et al., Relationship of interleukin-10 plasma levels to severity of injury and clinical outcome in injured patients, *J Trauma* 42, 863-870; discussion 870-861 (1997). [44] Lyons, A., Kelly, J. L., Rodrick, M. L., Mannick, J. A. & Lederer, J. A., Major injury induces increased production of interleukin-10 by cells of the immune system with a negative impact on resistance to infection, *Ann Surg* 226, 450-458; discussion 458-460 (1997). [45] Regan, M. C., *Host defense dysfunction in trauma, shock and sepsis: mechanisms and therapeutic approaches*, (Springer, Berlin, 1993). [46] Robson, M. C., Wound infection. A failure of wound healing caused by an imbalance of bacteria, *Surg Clin North Am* 77, 637-650 (1997). [47] Medzhitov, R., Recognition of microorganisms and activation of the immune response, *Nature* 449, 819-826 (2007). [48] Dinarello, C. A., Proinflammatory cytokines, *Chest* 118:503-508 (2000). [49] Foex, B. A., et al., Early cytokine response to multiple injury, *Injury* 24:373-376 (1993). [50] Jemal A, Siegel R, Ward E, et al., Cancer Statistics 2009, CA Cancer J Clin 2009 July-August; 59(4):225-49. [51] Cancer Facts and Figures 2008, Atlanta, Ga.; American Cancer Society: 2009. [52] Saving Women's Lives: Strategies for Improving Breast Cancer Detection and Diagnosis, Joy J E, Penhoet E E, and Petitti D B, eds., Institute of Medicine and National Research Council of the National Academies, National Academy Press, Washington D.C., 2004. [53] Winchester D P, Osteen R T, Menck H